

State: Georgia  
Grant Number: 08-953  
Study Number: 6

## LONG RANGE PERFORMANCE REPORT

Grant Title: State Funded Wildlife Survey

Period Covered: July 1, 2006 - June 30, 2007

Study Title: Wild Turkey Production and Population Indices

Study Objectives:

1. To determine annually an index of statewide turkey populations and production success in Georgia.
2. To organize data obtained in a form so that it can be used in sound management of turkeys in Georgia.

### Abstract

Recent analysis of long-term production data indicated that a new production index, 'Poults+Hens' instead of 'Poults/Observer' was the better predictor for Hours Hunted/Turkey Seen. Twenty-nine percent more Poults+Hens were observed in 2006 (5,787) versus 2005 (4,109) corresponding with the harvest season population index (Hours Hunted/Turkey Seen), which was 11% greater in 2006 (1.6) than 2005 (1.8). With the new analysis an inverse correlation coefficient of  $r = -0.90$  was obtained between the new production index and population indices for the entire survey period which began in 1978. Hunter success decreased slightly to 67.9% in 2007 from 69.1% in 2006. The average number of poults per hen was 1.6, which was up 6% from 2005.

#### A. Activity:

Job A. Turkey Production Index Survey - This survey was conducted during the months of May through August from 1978 to 1991. Beginning in 1991, the survey period was shortened to June through August when statistical analysis of data indicated the shorter time period was adequate. Data collection and summary for the 2007 survey period is not complete.

Cooperators involved in data collection for this survey were field personnel of the Game Management Section, Fisheries Management Section, and Law Enforcement Section of the Wildlife Resources Division. Observations were made during the course of regular field duties. No special efforts were made to locate turkeys for the survey.

Records were maintained of all turkey broods and hens, with and without broods. Data were compiled on a statewide and physiographic region basis. Historically, the

average number of poults seen per observer was the best index of production, however, recent analysis indicated this was not the case with data between 1987-2006. Currently, the best index of production data is estimated Total Poults+Hens.

Job B. Turkey Hunting Population Index Survey - The hunter cooperators participating in the survey were obtained from names of prospects submitted by WRD personnel and current cooperators. Cooperators were also solicited through newspaper and magazine requests and programs to interest groups. Randomly selected members of the Georgia Chapter of the National Wild Turkey Federation also were contacted to bring the total potential cooperating hunters to 2,000.

This survey is conducted during the regular spring gobbler-hunting season, which begins the first Saturday after March 19 and ends May 15. Specific information requested about each hunting trip was the date, hours hunted, county or physiographic region hunted, the number of turkeys seen, and the number of gobblers heard. Kill information was also requested, but was an optional item. Hunt record forms were supplied to all cooperators along with full instructions and a short newsletter on survey findings from previous years.

The number of turkeys observed per unit of hunting effort is used as an index of the hunting season population. The correlation between the population indices and the production indices are used in evaluating annual production and populations and in making comparisons for trends. Data were calculated on a statewide and physiographic region basis.

B. Target Date for Achievement and Accomplishments:

Job A. Planned dates and dates of accomplishment coincide, June 30, 2007.

Job B. Planned dates and dates of accomplishment coincide, June 30, 2007.

C. Significant Deviations:

Jobs A & B. None

D. Finds:

Job A. In 2006, 426 broods were observed (Table 1). This total is substantially more than in 2005, when 248 broods were observed and better than two years ago (2004) of 354 and close to what was observed three years ago (2003) of 448. The average brood size of 8.4 poults was 16% less than last year's average of 10.0. Twenty-nine percent more Poults+Hens were observed in 2006 (5,787) versus 2005 (4,109; Table 6).

Regional examination of poults/observer revealed that statewide it too was greater by 19% for 2006 (15.88) compared to 2005 (12.89; Table 3). Poults/observer was up

in all physiographic regions from 2005, except for the Lower Coastal Plain, which was down by 10%. The index for Ridge and Valley (RV or I) and Blue Ridge Mountains increased greater than 30% from 2005.

The number of hens reported totaled 2,208 (Table 4). The percent of hens with poults, 42% was 12% more than the 2005 total (Table 5). The average number of poults per hen, 1.6, increased by 6% from 2005 and therefore overall production was still considered poor for 2006. Historically, with Georgia's expanding turkey population an average of 3 poults per hen was considered good, however, recent data with a more stable population indicates that productivity threshold of  $\geq 2.0$  poults per hen may be an indicator of good reproductive levels.

Job B. Usable hunt data was supplied by 467 cooperators. Of these 404 came from the permanent cooperator list, 50 from the NWTF list, and 13 were new which resulted in a reporting rate (after deleting wrong addresses, deceased, quit hunting, incorrect data collection, etc.) of 38.2% and 6.2% from the permanent and NWTF list cooperators, respectively. These cooperators reported spending a total of 16,945.9 hours hunting (Table 7). The average season hunter effort was 10.5 trips totaling 36.3 hours. They reported observing 10,511 turkeys and hearing 8,083 gobblers. The statewide population index of 1.6 was 11% greater than last year (1.8 hrs hunter/turkey seen [the lower the number the greater the population]; Table 8). The effort per gobbler heard of 2.1 was 9% less than for the 2006 season (2.3; Table 8). The least hunting effort per turkey seen occurred in the Ridge and Valley and Lower Coastal Plain. The effort per gobbler heard was least in Upper and Lower Coastal Plain and highest in the Blue Ridge Mountains.

Statewide peak gobbling activity, 3.0 gobblers heard per trip, occurred during the first weekend (March 24-25). The next highest period was the fifth weekend (April 21-22) of the season with 1.9 gobblers heard per trip. All other periods averaged between 0.9 and 1.8 gobblers per trip, with the last weekend (May 14-15) being the lowest at 0.9 per trip. For most of the state the greatest amount of gobbling activity was the first 10 days (Mar 24 – April 1; Table 9). Other peaks in gobbling occurred during the fifth weekend (April 21-22) for Regions III, IV, and V. Even though the greatest amount of gobbling activity was the first weekend for most of the state, the greatest amount for Region II – Blue Ridge Mountains was the first week through the second weekend (Mar 26 – April 1; see tables 9 and 13).

The statewide gobbler harvest during the first seven days of the season amounted to 36% of the total season harvest, which was almost equal to 2006 (37%; Table 10). Peak harvest was generally seen within the first seven days of the season in all parts of the state (Tables 11 and 12).

Similar to previous seasons, the greatest number of trips were made during the first seven days of the season (Tables 13 and 14). Only minor variations in hunting effort measures have occurred over the years.

Hunter success decreased slightly to 67.9 % with 317 of 467 hunters reported taking or assisting in taking at least one gobbler (was 69.1% in 2006). Of the successful hunters, 115 (24.6 %) took or assisted in taking one bird, 87 (18.6 %) took or assisted in taking two birds, and 115 (24.6 %) took or assisted in taking three birds. Cooperators reported 194 gobblers killed by companions.

The predictive model analysis uses Poults+Hens of the reproductive season during the current year to predict the following years harvest season population index of Hours Hunted/Turkey Seen, where the predictor model (1978-2007) is:

**Constant + (Slope \*2006 Total Poults+Hens) = 2007 Hours Hunted/Turkey Seen**

Therefore:

$$3.3335 + (-0.00034*5,787) = 1.3 \text{ Hours Hunted/Turkey Seen in 2007.}$$

The predicted value of 1.3 was close to the actual observed value of 1.6 (2007). A relatively high inverse correlation  $r = -0.90$  was obtained from the comparison of the new nesting season population index versus the following years harvest season population index.

Table 1. Turkey broods and poult counts observed statewide in Georgia, 1978-2006.

Year	Broods		Poults	
	Total	Poult Counts	Brood Average	Est. Total
1978	123	82	8.6	1,058
1979	183	160	8.6	1,565
1980	176	169	8.4	1,479
1981	264	241	7.6	2,006
1982	260	218	7.7	2,002
1983	298	261	8.8	2,622
1984	293	247	6.8	1,992
1985	324	274	7.2	2,333
1986	430	377	9.4	4,042
1987	347	328	9.7	3,366
1988	347	321	7.9	2,741
1989	322	306	9.0	2,898
1990	459	278	7.6	3,488
1991	289	213	7.1	2,039
1992	298	274	6.8	2,027
1993	328	303	8.2	2,676
1994	341	316	9.4	3,209
1995	408	386	10.4	4,209
1996	271	239	7.5	2,033
1997	408	304	6.5	2,613
1998	595	534	7.0	4,185
1999	447	364	7.1	3,170
2000	393	358	7.2	2,809
2001	493	431	7.0	3,017
2002	648	618	6.0	3,894
2003	448	448	5.9	2,619
2004	354	354	10.6	3,733
2005	248	248	10.0	2,469
2006	426	426	8.4	3,579

Table 2. Turkey brood observations by physiographic region and month in Georgia, 2006.

Month	Region <sup>1</sup>					Total
	I	II	III	IV	V	
June	24	11	16	48	24	123
July	37	11	40	36	27	151
August	32	19	53	36	12	152
Totals	93	41	109	120	63	426

<sup>1</sup>Roman numerals correspond to physiographic regions as follows:

I - Valley and Ridge Lookout Mountain Plateau

II - Blue Ridge Mountains

III - Piedmont

IV - Upper Coastal Plain

V - Lower Coastal Plain

Table 3. Average number of turkey poults seen per observer (production index) in Georgia, 1978-2006.

Physiographic																
Region	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993
I	4.84	0	4.80	3.45	3.52	10.30	9.09	7.20	23.19	27.87	22.10	30.70	18.92	21.19	15.93	26.75
II	11.18	5.70	3.85	5.32	10.36	21.21	16.54	7.90	36.62	19.79	34.61	21.82	19.89	7.07	12.89	17.31
III	7.04	8.88	11.13	12.12	14.79	20.24	11.01	15.93	22.99	23.11	18.80	21.72	23.06	20.69	15.90	22.03
IV	3.86	5.16	5.23	7.15	11.44	9.42	8.78	15.03	23.03	11.54	12.01	12.72	10.83	7.71	7.84	14.91
V	6.28	7.36	3.63	8.89	5.37	5.19	6.37	10.93	13.74	6.60	9.32	8.12	20.10	5.27	10.32	11.15
Statewide	7.50	6.33	7.31	8.72	10.77	13.29	10.02	13.07	22.42	17.31	16.05	17.53	18.88	12.01	12.39	16.39

Table 3. Continued.

Physiographic													
Region	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
I	38.68	66.3	32.3	20.8	42.9	30.3	33.6	48.8	47.3	40.27	34.65	28.96	52.27
II	20.11	22.06	16.2	13.7	21.5	19.9	37.0	32.2	23.2	13.63	23.10	14.28	20.92
III	25.22	48.99	26.9	26.6	29.5	18.2	22.5	24.4	28.8	14.94	19.11	12.66	15.14
IV	19.17	21.0	16.5	14.1	22.6	21.2	17.4	18.9	21.7	8.55	16.18	12.10	14.62
V	8.00	14.83	4.5	9.1	6.2	11.0	8.1	9.6	13.9	10.86	13.42	10.36	9.29
Statewide	20.63	31.78	18.9	16.2	22.1	17.7	18.2	21.3	24.1	13.11	18.28	12.89	15.88

Table 4. Turkey hens observed with poults, without poults, and uncertain of accompanying poults statewide in Georgia, 1978-2006.

Year	Hens Reported			Total
	With Poults	Without Poults	Uncertain of Poults	
1978	145	70	26	241
1979	176	131	39	346
1980	166	133	15	314
1981	276	116	66	458
1982	327	136	24	487
1983	361	211	72	644
1984	261	232	59	552
1985	475	251	81	807
1986	648	283	84	1,015
1987	519	230	52	801
1988	529	305	59	893
1989	459	261	48	768
1990	642	371	49	1,062
1991	321	399	59	779
1992	407	490	59	956
1993	374	292	41	707
1994	463	361	66	890
1995	606	301	83	990
1996	298	384	74	756
1997	560	618	271	1,449
1998	820	661	236	1,717
1999	560	753	344	1,657
2000	734	577	251	1,562
2001	634	589	337	1,560
2002	695	644	220	1,559
2003	795	1,113	296	2,204
2004	930	586	347	1,863
2005	611	772	257	1,640
2006	932	864	412	2,208



Table 5. Percent of turkey hens accompanied by poults (2nd potential population index) and the average number of poults per hen statewide in Georgia, 1978-2006.

Year	Percent Hens With Poults	Poults Per Hen
1978	60	4.4
1979	51	4.5
1980	53	4.7
1981	60	4.4
1982	67	4.1
1983	56	4.1
1984	47	3.6
1985	59	3.6
1986	64	4.4
1987	65	4.2
1988	59	3.1
1989	60	3.8
1990	60	3.3
1991	41	2.6
1992	43	2.1
1993	56	3.8
1994	56	3.6
1995	61	4.3
1996	39	2.7
1997	39	1.8
1998	48	2.4
1999	34	1.9
2000	47	1.8
2001	41	2.2
2002	45	2.5
2003	36	1.2
2004	50	2.0
2005	37	1.5
2006	42	1.6

Table 6. Estimated Total Poults + hens population indices in Georgia, 1978-2006.

Population Index	Nesting Season	Statewide Poults+Hens
	1978	1,299
	1979	1,911
	1980	1,793
	1981	2,464
	1982	2,489
	1983	3,266
	1984	2,544
	1985	3,140
	1986	5,057
	1987	4,167
	1988	3,634
	1989	3,666
	1990	4,550
	1991	2,758
	1992	2,983
	1993	3,383
	1994	4,099
	1995	5,199
	1996	2,789
	1997	4,062
	1998	5,902
	1999	4,827
	2000	4,371
	2001	4,577
	2002	5,453
	2003	4,823
	2004	5,596
	2005	4,109
	2006	5,787

Table 7. Summary of turkey hunter cooperator data in Georgia, 2007.

Item	Physiographic Region <sup>1</sup>					Statewide
	I	II	III	IV	V	
Total Hunters	58	35	275	181	54	467**
Total Hours	1,617	634	8,312.95	4,989.2	1,392.75	16,945.9
Total Trips	484	163	2,192	1,576	489	4,904
Avg. Hours	27.9	18.1	30.2	27.6	25.8	36.3
Avg. Trips	8.3	4.7	8.0	8.7	9.1	10.5
Avg. Hrs./Trip	3.3	3.9	3.8	3.2	2.8	3.5
Total Turkeys Seen	1,353	400	4,101	3,310	1,347	10,511
Hrs./Turkeys Seen	1.2	1.6	2.0	1.5	1.0	1.6
Total Gobblers Heard	694	148	3,435	2,927	879	8,083
Hrs./Gobbler Heard	2.3	4.3	2.4	1.7	1.6	2.1
Total Kill*	60	19	278	281	96	734
Companion Killed	16	6	53	92	27	194
Hours/Kill	27.0	33.4	29.9	17.8	14.5	23.1

<sup>1</sup>Roman numerals correspond to physiographic regions as follows:

- I - Ridge and Valley
- II - Blue Ridge Mountains
- III - Piedmont
- IV - Upper Coastal Plain
- V - Lower Coastal Plain

\*includes both gobblers taken and assisted in taking

\*\* less than Regions summed because some hunters hunted in more than one Region

Table 8. Turkey hunting population indices in Georgia, 1979-2007.

Population Index	Hunt Season	Physiographic Region					Statewide
		I	II	III	IV	V	
Hours/Turkey Seen	1979	20.5	3.5	2.9	3.1	2.8	3.0
	1980	1.6	6.0	2.9	2.6	2.4	3.1
	1981	1.5	4.7	2.2	3.2	2.8	2.5
	1982	2.2	5.0	2.8	3.3	1.8	2.9
	1983	2.5	3.1	2.2	2.0	1.9	2.3
	1984	2.2	4.1	2.4	1.6	1.5	2.3
	1985	2.3	3.4	2.6	2.5	3.5	2.6
	1986	3.2	4.6	2.3	2.0	3.4	2.5
	1987	4.1	2.9	2.6	1.7	2.1	2.4
	1988	1.0	2.9	1.9	1.6	2.1	1.8
	1989	1.7	2.3	2.3	1.6	1.2	1.9
	1990	1.8	2.8	2.0	1.9	1.7	2.0
	1991	1.6	2.3	2.0	1.7	1.8	1.9
	1992	1.4	2.7	2.4	1.7	2.3	2.1
	1993	2.0	4.0	2.5	1.6	1.6	2.1
	1994	2.4	2.2	2.1	1.6	1.4	1.9
	1995	1.7	2.2	2.4	1.8	2.0	2.1
	1996	1.2	1.8	1.6	1.6	1.5	1.5
	1997	1.0	2.1	1.8	1.5	1.3	1.6
	1998	1.0	1.9	1.9	1.7	1.4	1.7
	1999	0.9	2.7	1.5	1.4	1.5	1.4
	2000	1.4	2.3	2.0	1.5	1.5	1.7
	2001	4.2	3.4	1.3	1.7	1.4	1.7
	2002	3.9	3.7	1.2	2.2	1.9	2.6
	2003	1.5	1.8	1.6	1.4	1.5	1.5
	2004	1.1	2.2	1.7	1.2	1.3	1.4
	2005	1.1	2.7	2.2	1.4	1.2	1.6
	2006	1.2	2.0	2.3	1.6	1.2	1.8
2007	1.2	1.6	2.0	1.5	1.0	1.6	

Table 8. Continued.

Population Index	Hunt Season	Physiographic Region					Statewide
		I	II	III	IV	V	
Hours/Gobbler Heard	1979	50.7	7.3	3.3	2.1	1.8	3.2
	1980	2.9	4.7	3.4	2.9	9.1	3.4
	1981	2.9	4.4	3.0	2.3	2.0	2.9
	1982	3.1	3.6	3.0	2.3	2.3	2.9
	1983	4.4	2.8	3.3	2.0	2.4	2.8
	1984	3.1	5.2	3.3	1.8	1.4	3.0
	1985	2.4	4.2	2.9	1.8	3.0	2.6
	1986	2.6	3.4	2.1	1.3	1.6	2.0
	1987	2.2	5.2	2.4	1.7	2.0	2.4
	1988	1.5	2.6	2.7	1.4	1.6	2.2
	1989	2.1	2.1	2.1	1.5	2.1	1.9
	1990	2.3	4.2	2.5	1.7	1.7	2.2
	1991	2.7	5.5	2.7	2.0	2.9	2.7
	1992	2.4	4.2	2.9	1.8	1.6	2.6
	1993	3.2	6.3	3.6	2.1	2.7	3.1
	1994	3.4	6.1	3.5	1.9	2.2	2.9
	1995	2.0	3.3	2.5	1.9	2.1	2.3
	1996	3.3	3.5	2.7	2.0	2.1	2.5
	1997	2.3	5.6	2.2	1.6	2.2	2.2
	1998	2.5	4.1	2.7	1.9	2.1	2.4
	1999	2.7	3.7	2.8	1.7	2.0	2.4
	2000	2.1	3.8	2.2	1.8	1.8	2.1
	2001	4.8	5.4	1.8	2.4	2.7	2.4
	2002	4.2	4.9	1.6	2.8	2.6	3.2
	2003	1.9	2.0	1.8	2.1	1.8	1.9
	2004	2.0	4.2	2.4	1.6	1.7	2.0
	2005	2.5	4.3	2.9	1.8	1.9	2.4
	2006	2.2	3.2	2.7	1.9	1.7	2.3
2007	2.3	4.3	2.4	1.7	1.6	2.1	

Table 8. Continued.

Population Index	Hunt Season	Physiographic Region					Statewide
		I	II	III	IV	V	
Hours/Gobbler Killed	1979	96.5	79.8	35.1	27.5	23.3	35.7
	1980	13.2	35.7	39.6	35.8	19.1	35.9
	1981	10.7	29.5	31.0	29.9	23.0	30.7
	1982	25.5	90.3	29.7	30.0	19.0	31.3
	1983	30.9	29.7	27.8	28.3	22.6	27.4
	1984	31.1	45.8	35.3	31.4	12.8	34.0
	1985	22.2	48.2	38.7	24.0	32.4	33.6
	1986	23.0	42.1	28.6	21.9	16.0	26.7
	1987	35.4	68.3	30.4	25.8	32.1	32.1
	1988	17.6	25.3	35.9	18.9	18.7	28.0
	1989	22.6	41.4	29.8	17.0	21.1	24.8
	1990	29.8	55.2	29.3	26.4	16.3	28.3
	1991	42.7	48.4	36.9	24.7	23.2	33.9
	1992	44.9	49.4	45.3	20.9	22.0	36.7
	1993	32.2	46.5	46.0	19.8	38.7	34.9
	1994	36.2	42.0	36.9	20.9	18.7	30.1
	1995	25.4	29.9	25.3	18.6	18.7	22.7
	1996	28.9	34.1	29.3	25.9	26.0	26.8
	1997	28.7	38.8	31.9	19.6	20.7	27.7
	1998	29.2	35.8	29.2	23.3	19.0	26.3
1999	28.0	50.6	33.6	19.1	24.2	27.8	
2000	27.8	34.0	28.5	22.9	23.0	26.4	
2001	60.6	48.3	22.6	25.7	23.2	27.9	
2002	59.7	43.6	21.1	27.6	19.2	34.2	
2003	21.6	22.8	26.7	26.4	25.4	25.7	
2004	21.5	44.6	27.4	18.5	21.2	23.4	
2005	26.3	42.3	31.0	18.0	18.1	24.4	
2006	20.8	40.2	31.0	21.6	16.9	25.1	
2007	27.0	33.4	29.9	17.8	14.5	23.1	

Table 9. Number of gobblers heard per hunting trip in Georgia, 2007.

Date		Physiographic Region					Statewide
Weekend	Weekday	I	II	III	IV	V	
3/24-3/25		2.7	0.8	2.8	3.5	3.4	3.0
	3/26-3/30	1.4	1.4	1.8	2.0	2.0	1.8
3/31-4/01		1.9	1.3	1.6	2.1	2.0	1.8
	4/02-4/06	1.2	1.1	1.2	1.3	2.0	1.3
4/07-4/08		0.7	0.0	1.0	1.4	0.9	1.1
	4/09-4/13	1.2	1.3	1.5	1.7	1.6	1.5
4/14-4/15		1.1	0.6	1.5	1.9	1.9	1.6
	4/16-4/20	1.3	1.3	1.1	1.4	1.1	1.2
4/21-4/22		1.3	0.5	1.8	2.2	2.2	1.9
	4/23-4/27	1.1	0.6	1.4	1.3	1.6	1.3
4/28-4/29		1.6	0.6	1.3	1.7	1.5	1.4
	4/30-5/04	1.6	0.4	1.0	1.8	1.4	1.4
5/05-5/06		1.1	1.0	0.9	1.1	1.1	1.0
	5/07-5/11	1.2	0.9	0.9	1.0	1.1	1.0
5/12-5/13		0.8	0.3	0.9	1.2	1.6	1.0
	5/14-5/15	1.0	0.8	0.8	1.2	0.5	0.9
Season		1.4	0.9	1.6	1.9	1.8	1.6

Table 10. Chronological summary of gobbler harvest in Georgia, 2007.

Date		Gobblers	% of Season Kill*	
Weekend	Weekday	Killed	Date	Cumulative
3/24-3/25		154	21	21
	3/26-3/30	112	15	36
3/31-4/01		64	9	45
	4/02-4/06	80	11	56
4/07-4/08		21	3	59
	4/09-4/13	47	6	65
4/14-4/15		34	5	70
	4/16-4/20	42	6	76
4/21-4/22		28	4	80
	4/23-4/27	36	5	85
4/28-4/29		29	4	89
	4/30-5/04	24	3	92
5/05-5/06		19	3	95
	5/07-5/11	20	3	98
5/12-5/13		16	2	100
	5/14-5/15	8	1	101
Total		734	101	101

\*over 100% because of rounding

Table 11. Chronological distribution of gobbler harvest by physiographic region in Georgia, 2007.

Dates		Physiographic Region					Statewide
Weekend	Weekday	I	II	III	IV	V	
3/24-3/25		12	3	60	66	13	154
	3/26-3/30	8	3	60	25	16	112
3/31-4/01		4	3	21	29	7	64
	4/02-4/06	3	1	32	32	12	80
4/07-4/08		4	0	3	9	5	21
	4/09-4/13	3	2	14	21	7	47
4/14-4/15		4	0	16	10	4	34
	4/16-4/20	6	2	16	13	5	42
4/21-4/22		1	1	9	15	2	28
	4/23-4/27	3	0	10	12	11	36
4/28-4/29		5	0	9	13	2	29
	4/30-5/04	2	0	5	10	7	24
5/05-5/06		2	1	7	7	2	19
	5/07-5/11	2	0	6	11	1	20
5/12-5/13		0	0	8	6	2	16
	5/14-5/15	1	3	2	2	0	8
Season		60	19	278	281	96	734

Table 12. Chronological distribution of gobbler harvest (%) by physiographic region in Georgia, 2007.

Date		Physiographic Region					Statewide
Weekend	Weekday	I	II	III	IV	V	
3/24-3/25		20	16	22	23	14	21
	3/26-3/30	13	16	22	9	17	15
3/30-4/01		7	16	8	10	7	9
	4/02-4/06	5	5	12	11	13	11
4/07-4/08		7	0	1	3	5	3
	4/09-4/13	5	11	5	7	7	6
4/14-4/15		7	0	6	4	4	5
	4/16-4/20	10	11	6	5	5	6
4/21-4/22		2	5	3	5	2	4
	4/23-4/27	5	0	4	4	11	5
4/28-4/29		8	0	3	5	2	4
	4/30-5/04	3	0	2	4	7	3
5/05-5/06		3	5	3	2	2	3
	5/07-5/11	3	0	2	4	1	3
5/12-5/13		0	0	3	2	2	2
	5/14-5/15	2	16	1	1	0	1



Table 13. Chronological distribution of turkey hunting trips by physiographic region in Georgia, 2007.

Dates		Physiographic Region					Statewide
Weekend	Weekday	I	II	III	IV	V	
3/24-3/25		52	12	302	191	49	605
	3/26-3/30	56	15	310	196	59	637
3/31-4/01		34	11	187	124	39	395
	4/02-4/06	42	19	205	171	53	490
4/07-4/08		25	1	88	71	28	213
	4/09-4/13	31	15	162	110	47	365
4/14-4/15		21	9	111	72	22	235
	4/16-4/20	46	17	133	104	36	336
4/21-4/22		28	8	118	85	21	260
	4/23-4/27	29	11	134	114	41	329
4/28-4/29		23	9	103	73	19	227
	4/30-5/04	31	9	92	72	24	228
5/05-5/06		16	3	64	53	17	153
	5/07-5/11	28	8	74	68	16	194
5/12-5/13		8	11	72	41	12	144
	5/14-5/15	14	5	37	31	6	93
Season		484	163	2,192	1,576	489	4,904

Table 14. Chronological distribution of turkey hunting trips (%) by physiographic region in Georgia, 2007.

Dates		Physiographic Region					Statewide
Weekend	Weekday	I	II	III	IV	V	
3/24-3/25		11	7	14	12	10	12
	3/26-3/30	12	9	14	12	12	13
3/31-4/01		7	7	9	8	8	8
	4/02-4/06	9	12	9	11	11	10
4/07-4/08		5	1	4	5	6	4
	4/09-4/13	6	9	7	7	10	7
4/14-4/15		4	6	5	5	4	5
	4/16-4/20	10	10	6	7	7	7
4/21-4/22		6	5	5	5	4	5
	4/23-4/27	6	7	6	7	8	7
4/28-4/29		5	6	5	5	4	5
	4/30-5/04	6	6	4	5	5	5
5/05-5/06		3	2	3	3	3	3
	5/07-5/11	6	5	3	4	3	4
5/12-5/13		2	7	3	3	2	3
	5/14-5/15	3	3	2	2	1	2